arXiv:1704.00837v1 [math.CA] 4 Apr 2017

A NOTE ON BAND-LIMITED MINORANTS OF AN EUCLIDEAN BALL

FELIPE GONÇALVES

Abstract. These are notes on the Beurling-Selberg problem of finding band-limited L^1-functions that lie below the indicator function of an Euclidean ball. We compute the critical radius of the support of the Fourier transform for which such construction is non-trivial.

1. Introduction

For a given $r > 0$ we denote by $B^d(r)$ the closed Euclidean ball in $\mathbb{R}^d$ centered at the origin with radius $r > 0$. We simply write B^d when $r = 1$. Define the following quantity

$$\beta(d, r) = \sup_F \int_{\mathbb{R}^d} F(\boldsymbol{x}) dx, \tag{1.1}$$

where the supremum is taken among functions $F \in L^1(\mathbb{R}^d)$ such that:

(1) The Fourier Transform of $F(x)$,

$$\widehat{F}(\xi) = \int_{\mathbb{R}^d} F(x) e^{2\pi i x \cdot \xi} \mathrm{d}x,$$

is supported in $B^d(r)$;

(2) $F(x) \leq \mathbf{1}_{B^d}(x)$ for all $x \in \mathbb{R}^d$.

We call such a function $\beta(d, r)$-admissible. A trivial observation is that $F \equiv 0$ is $\beta(d, r)$-admissible, hence $\beta(d, r) \geq 0$. Heuristically, such function $F(x)$ should exist and its mass should be close to $\mathrm{vol}(B^d)$ when r is large. On the other hand, if r is small, the mass of $F(x)$ should be close to zero and a critical $r_d > 0$ should exist such that no function can beat the identically zero function for $r \leq r_d$. For this reason we define

$$r_d = \inf\{r > 0 : \beta(d, r) > 0\}$$

and it is this critical radius that we want to study in this paper.

The problem stated in (1.1) has its origins with Beurling and Selberg which studied one-sided band-limited approximations for many different functions other than indicator functions with the purpose of using them to derive sharp estimates in analytic number theory (see the introduction of [6] for a nice first view). Although

Date: April 5, 2017.

Selberg was one of the first to study the higher dimensional problem, it was first systematically analysed by Holt and Vaaler in the remarkable paper [4]. They were able to construct non-zero $\beta(N,r)$-admissible functions for any $r>0$ and, most importantly, they established a fascinating connection of the d-dimensional problem with the theory of Hilbert spaces of entire functions contructed by de Branges (see [1]). They reduced the higher dimensional problem, after a radialization argument, to a weighted one-dimensional problem where the weight was given by a special function of Hermite-Biehler class, which in turn allowed them to use the machinery of homogeneous de Branges spaces to attack the problem. This new connection established by Holt and Vaaler started a new way of thinking about these kind of problems and ultimately inspired Littmann to completely solve the one-dimensional problem in [5] by using a cleaver argument based on a special structure of certain de Branges spaces. Finally, using the ideas introduced by Littmann in [5], the problem of minorizing the indicator function of a symmetric interval was completely solved in [2] in the de Branges space setting[1].

1.1. **Main Result.** For any given parameter $\nu>-1$ let $J_\nu(z)$ denote the classical Bessel function of the first kind. We also denote by $\{j_{\nu,n}\}_{n\geq 1}$ its positive zeros listed in increasing order. The following is the main result of this paper.

Theorem 1. *We have*

$$r_d=\frac{j_{d/2-1,1}}{\pi}.$$

Moreover, if $j_{d/2-1,1}<\pi r<j_{d/2,1}$ *then*

$$\beta(d,r)=\frac{(2/r)^d}{|\mathbb{S}^d|}\frac{\gamma_{\pi r}}{1+\gamma_{\pi r}/d},$$

where $\gamma_{\pi r}=-\frac{\pi r J_{d/2-1}(\pi r)}{J_{d/2}(\pi r)}>0$. *In particular we have*

$$\beta(d,r)=\frac{\pi^2 2^d}{r^{d-1}|\mathbb{S}^d|}\left(r-\frac{j_{d/2-1,1}}{\pi}\right)+O_d\left(r-\frac{j_{d/2-1,1}}{\pi}\right)^2$$

for r *close to* $\frac{j_{d/2-1,1}}{\pi}$.

Remarks.

(1) It is known that $j_{\nu,1}=\nu+1.855757\nu^{1/3}+O(\nu^{-1/3})$ as $\nu\to\infty$ (see [3, Section 1.3]). This implies that $r_d=\frac{d}{2\pi}+\frac{1.855757}{2^{1/3}\pi}d^{1/3}+O(d^{-1/3})$ as $d\to\infty$. Heuristically, this means that if one wishes to non-trivially minorate (that is, beat the zero function) the indicator function of a ball of radius of order $\sqrt{d}$ then one needs frequencies of order at least $\sqrt{d}$. The first 5 values of r_d rounded up to 4 significant digits are the following: $r_1=1/2$, $r_2=0.7655$, $r_3=1$, $r_4=1.220$ and $r_5=1.431$.

[1]It is not the intention of this paper to give a survey of related articles on the subject, which is very rich and full of subtleties, the purpose here is to drawn a strait line between what he have so far and what we want to show.

(2) Explicit expressions for $\beta(d,r)$ can also be tracked from [2, Theorem 5], but they involve sums of Bessel functions evaluated at Bessel zeros that can be quite complicated to grasp. Moreover, this is the case only when πr is a zero of $J_{d/2-1}(z)$ or $J_{d/2}(z)$. If that is not the case, then writing a formula for $\beta(d,r)$ becomes pointless, since it will involve zeros of more complicated functions related to Bessel functions and this is not the purpose here.

2. Proof of Theorem 1

Step 1. The first step is to reduce the higher dimensional by considering only radial functions. We can apply [4, Lemmas 18 and 19] to reduce the d-dimensional problem to the following weighted one-dimensional problem

$$\beta(d,r) = \frac{|\mathbb{S}^{d-1}|}{2} \sup_F \int_{\mathbb{R}} F(x)|x|^{d-1}dx, \tag{2.1}$$

where $|\mathbb{S}^{d-1}|$ denotes the surface area of the unit sphere in $\mathbb{R}^d$ and the supremum is taken among functions $F \in L^1(\mathbb{R}, |x|^{d-1}\mathrm{d}x)$ such that:

(1) $F(x)$ is the restriction to the real axis of an even entire function $F(z)$ of exponential type at most $2\pi r$, that is,

$$|F(z)| \leq Ce^{2\pi r|\operatorname{Im} z|}, \quad z \in \mathbb{C}$$

for some constant $C > 0$;

(2) $F(x) \leq \mathbf{1}_{[-1,1]}(x)$ for all $x \in \mathbb{R}$.

This problem in such framework becomes treatable with the theory de Branges spaces of entire functions. The latter generalize the well known Paley-Wiener spaces by using weighted norms given by Hermite-Biehler functions. In what follows we briefly review the construction of a special family of de Branges spaces called *homogeneous spaces* which were introduced by de Branges (see [1, Section 50] and [4, Section 5]). We refer to [4, Section 3] for a brief description of the general theory and also to [1, Chapter 2] for the full theory.

Step 2. Let $\nu > -1$ be a parameter and consider the real entire functions $A_\nu(z)$ and $B_\nu(z)$ given by

$$A_\nu(z) = \sum_{n=0}^{\infty} \frac{(-1)^n \left(\frac{1}{2}z\right)^{2n}}{n!(\nu+1)(\nu+2)\dots(\nu+n)} = \Gamma(\nu+1)\left(\tfrac{1}{2}z\right)^{-\nu} J_\nu(z)$$

and

$$B_\nu(z) = \sum_{n=0}^{\infty} \frac{(-1)^n \left(\frac{1}{2}z\right)^{2n+1}}{n!(\nu+1)(\nu+2)\dots(\nu+n+1)} = \Gamma(\nu+1)\left(\tfrac{1}{2}z\right)^{-\nu} J_{\nu+1}(z).$$

If we write

$$E_\nu(z) = A_\nu(z) - iB_\nu(z),$$

then $E_\nu(z)$ is a Hermite–Biehler function, that is, it satisfies the following fundamental inequality

$$|E_\nu(\overline{z})| < |E_\nu(z)|$$

for all $z \in \mathbb{C}$ with $\operatorname{Im} z > 0$. It also known that this function does not have real zeros, that $E(iy) \in \mathbb{R}$ for all real y (that is, $B_\nu(z)$ is odd and $A_\nu(z)$ is even), that $E_\nu(z)$ is of bounded type in the upper-half plane (that is, $\log|E_\nu(z)|$ has a positive harmonic majorant in the upper-half plane) and $E_\nu(z)$ is of exponential type 1. We also have that

$$c|x|^{2\nu+1} \leq |E_\nu(x)|^{-2} \leq C|x|^{2\nu+1}$$

for all $|x| \geq 1$ and for some $c, C > 0$. The homogeneous space $\mathcal{H}(E_\nu)$ is then defined as the space of entire functions $F(z)$ of exponential type at most 1 and such that[2]

$$\int_{\mathbb{R}} |F(x)|^2 \, |E_\nu(x)|^{-2} \, dx < \infty.$$

Using standard asymptotic expansions for Bessel functions one can show that $A_\nu, B_\nu \notin \mathcal{H}(E_\nu)$. As a particular case, observing that $E_{-1/2} = e^{-iz}$ we can deduce that $\mathcal{H}(E_{-1/2})$ coincides with the Paley-Wiener space of square integrable entire functions of exponential type at most 1.

These spaces are relevant to our problem since we have the following magical identity

$$a_\nu \int_{\mathbb{R}} |F(x)|^2 \, |E_\nu(x)|^{-2} \, \mathrm{d}x = \int_{\mathbb{R}} |F(x)|^2 \, |x|^{2\nu+1} \, \mathrm{d}x \tag{2.2}$$

for each $F \in \mathcal{H}(E_\nu)$, where $a_\nu = 2^{2\nu+1}\,\Gamma(\nu+1)^2/\pi$. Identity (2.2) also extends to the L^1-setting for functions which do not change in sign at infinity. Let $F(z)$ be an entire function of exponential type at most 2 such that $F(x) \leq \mathbf{1}_{[-t,t]})(x)$ for some $t > 0$ and $F \in L^1(\mathbb{R}, |x|^{2\nu+1}\mathrm{d}x)$. Since $G_n(x) = (\frac{\sin(x/n)}{x/n})^n$ belongs to $\mathcal{H}(E_\nu)$ for large n and it converges to 1 uniformly in compact sets as $n \to \infty$, we have that $4G_n(x)^2 - F(x) \geq 0$ for all real x (if n is large and even) and this function has exponential type at most 2. This implies that $4G_n(x)^2 - F(x) = H_n(z)\overline{H_n(\overline{z})}$ for all $z \in \mathbb{C}$, for some entire function $H_n(z)$ of exponential type at most 1 (see [1, Theorem 13]). We conclude that $H_n \in \mathcal{H}(E_\nu)$ and we obtain

$$\begin{aligned} a_\nu \int_{\mathbb{R}} F(x) \, |E_\nu(x)|^{-2} \mathrm{d}x &= a_\nu \int_{\mathbb{R}} \{|H_n(x)|^2 + 4G_n(x)^2\} \, |E_\nu(x)|^{-2} \mathrm{d}x \\ &= \int_{\mathbb{R}} \{|H_n(x)|^2 + 4G_n(x)^2\} |x|^{2\nu+1} \mathrm{d}x \\ &= \int_{\mathbb{R}} F(x) |x|^{2\nu+1} \mathrm{d}x. \end{aligned} \tag{2.3}$$

[2]As an historical note, de Branges originally defined this space in another way but, in [4, Lemma 16], they showed that this is an equivalent definition.

Step 3. Taking $\nu = d/2 - 1$, we can re-scale the type condition and use identities (2.3) and (2.1) to reduce the problem of minorizing the indicator function of an Euclidean ball to the following final form

$$\beta(d,r) = \frac{(2/r)^d}{\pi|\mathbb{S}^d|}\Lambda^-_{E_{d/2-1}}(\pi r),$$

where

$$\Lambda^-_{E_{d/2-1}}(\pi r) = \sup_F \int_{\mathbb{R}} F(x)|E_{d/2-1}(x)|^{-2}\mathrm{d}x$$

and the supremum is taken among functions $F \in L^1(\mathbb{R}, |E_{d/2-1}(x)|^{-2}\mathrm{d}x)$ such that:

(1) $F(x)$ is the restriction to the real axis of an even entire function $F(z)$ of exponential type at most 2;
(2) $F(x) \leq \mathbf{1}_{[-\pi r, \pi r]}(x)$ for all $x \in \mathbb{R}$.

The above problem was completely solved in [2]. By all the previous discussion in Step 2, we can apply [2, Theorem 5 (i) and (iv)] to the function $E_{d/2-1}(z)$ (it actually can be applied to any $E_\nu(z)$) to derive that $\pi r_d = j_{d/2-1,1}$. Moreover, if $j_{d/2-1,1} < \pi r < j_{d/2,1}$ then [2, Theorem 5 (iv)] also give us that

$$\Lambda^-_{E_{d/2-1}}(\pi r) = \frac{\pi\gamma_{\pi r}}{1+\gamma_{\pi r}/d},$$

where $\gamma_{\pi r} = -\frac{\pi r J_{d/2-1}(\pi r)}{J_{d/2}(\pi r)} > 0$. A simple Taylor expansion leads to

$$\Lambda^-_{E_{d/2-1}}(\pi r) = \pi^2 r(\pi r - j_{d/2-1,1}) + O(\pi r - j_{d/2-1,1})^2$$

and we finally obtain that

$$\beta(d,r) = \frac{\pi^2 2^d}{r^{d-1}|\mathbb{S}^d|}\left(r - \frac{j_{d/2-1,1}}{\pi}\right) + O_d\left(r - \frac{j_{d/2-1,1}}{\pi}\right)^2.$$

References

[1] L. de Branges, *Hilbert Spaces of Entire Functions*, Prentice-Hall Series in Modern Analysis, 1968.

[2] E. Carneiro, V. Chandee, F. Littmann and M. B. Milinovich, Hilbert spaces and the pair correlation of zeros of the Riemann zeta-function, (to appear in J. Reine Angew. Math).

[3] A. Elbert, Some recent results on the zeros of Bessel functions and orthogonal polynomials, Journal of Computational and Applied Mathematics 133 (2001), 65-83.

[4] J. J. Holt and J. D. Vaaler, The Beurling-Selberg extremal functions for a ball in Euclidean space, Duke Math. J., Volume 83, Number 1 (1996), 203-248.

[5] F. Littmann, Quadrature and extremal bandlimited functions, SIAM J. Math. Anal. 45 (2013), no. 2, 732-747.

[6] J. D. Vaaler, Some extremal functions in Fourier analysis, Bulletin of American Mathematical Society 12 (1985), no. 2, 183-216.

University of Alberta, Mathematical and Statistical Sciences, CAB 632, Edmonton, Alberta, Canada T6G 2G1

E-mail address: felipe.goncalves@ualberta.ca

URL: sites.ualberta.ca/~goncalve